



**Queensland University of Technology**  
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Smith, Graham (2014) Poly[ $\mu_5$ -{hydrogen bis[(E)-cinnamato]}]-caesium. *Acta Crystallographica. Section E: Structure Reports Online*, 70(2), m43-m44.

This file was downloaded from: <http://eprints.qut.edu.au/70898/>

**© Copyright 2014 The author**

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.

**Notice:** *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1107/S1600536814000804>

Acta Crystallographica Section E

Structure Reports


Online

ISSN 1600-5368

# Poly[ $\mu_5$ -{hydrogen bis[(*E*)-cinnamato}]-caesium]

**Graham Smith***Acta Cryst.* (2014). **E70**, m43–m44

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The average publication time is less than one month. Articles are published in a short-format style with enhanced supplementary materials. Each publication consists of a complete package – the published article, HTML and PDF supplements, CIF, structure factors, graphics, and any other submitted supplementary files.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

# Poly[ $\mu_5$ -{hydrogen bis[(*E*)-cinnamato]}-caesium]

Graham Smith

Science and Engineering Faculty, Queensland University of Technology, GPO Box 2434, Brisbane, Queensland 4001, Australia  
Correspondence e-mail: g.smith@qut.edu.au


Received 8 January 2014; accepted 13 January 2014

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(\text{C}-\text{C}) = 0.017$  Å;  $R$  factor = 0.071;  $wR$  factor = 0.144; data-to-parameter ratio = 16.0.

In the structure of the title polymeric complex,  $[\text{Cs}(\text{C}_9\text{H}_7\text{O}_2)(\text{C}_9\text{H}_8\text{O}_2)]_n$ , a caesium salt of *trans*-cinnamic acid, the  $\text{Cs}^+$  ions of the two individual irregular  $\text{CsO}_8$  coordination polyhedra lie on twofold rotation axes and are linked by four bridging carboxyl O-atom donors from two cinnamate ligand species. These two ligand components are interlinked through a delocalized H atom within a short  $\text{O} \cdots \text{H} \cdots \text{O}$  hydrogen bond. Structure extension gives a two-dimensional coordination polymer which lies parallel to (001). The structure was determined from a crystal twinned by non-merohedry, with a twin component ratio of approximately 1:1.

## Related literature

For the structures of the ammonium salts of hydrogen bis(3-chlorocinnamate) and hydrogen bis(3-bromocinnamate), see: Chowdhury & Kariuki (2006). For structures of alkali metal salts of ring-substituted *trans*-cinnamic acid, see: Kariuki *et al.* (1994, 1995); Crowther *et al.* (2008); Smith & Wermuth (2009, 2011). For the structure of *trans*-cinnamic acid, see: Wierda *et al.* (1989); Abdelmoty *et al.* (2005).



## Experimental

### Crystal data

$[\text{Cs}(\text{C}_9\text{H}_7\text{O}_2)(\text{C}_9\text{H}_8\text{O}_2)]$   
 $M_r = 428.21$   
Monoclinic,  $P2_1/c$   
 $a = 7.8608$  (6) Å  
 $b = 5.6985$  (7) Å  
 $c = 38.817$  (3) Å  
 $\beta = 98.733$  (6)°

$V = 1718.6$  (3) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 2.17$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.35 \times 0.35 \times 0.06$  mm

### Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer  
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2013)  
 $T_{\min} = 0.711$ ,  $T_{\max} = 0.980$

6675 measured reflections  
3353 independent reflections  
2552 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$   
 $wR(F^2) = 0.144$   
 $S = 1.19$   
3353 reflections

210 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.26$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -2.19$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

|                                 |           |                                |           |
|---------------------------------|-----------|--------------------------------|-----------|
| Cs1—O13 <i>B</i>                | 3.060 (8) | Cs2—O13 <i>B</i>               | 3.063 (8) |
| Cs1—O14 <i>A</i>                | 3.182 (8) | Cs2—O14 <i>A</i>               | 3.377 (9) |
| Cs1—O13 <i>A</i> <sup>i</sup>   | 3.132 (9) | Cs2—O13 <i>A</i> <sup>i</sup>  | 3.108 (9) |
| Cs1—O14 <i>B</i> <sup>i</sup>   | 3.183 (9) | Cs2—O14 <i>B</i> <sup>i</sup>  | 3.130 (9) |
| Cs1—O13 <i>B</i> <sup>ii</sup>  | 3.060 (8) | Cs2—O13 <i>B</i> <sup>iv</sup> | 3.063 (8) |
| Cs1—O14 <i>A</i> <sup>ii</sup>  | 3.182 (8) | Cs2—O14 <i>A</i> <sup>iv</sup> | 3.377 (9) |
| Cs1—O13 <i>A</i> <sup>iii</sup> | 3.132 (9) | Cs2—O13 <i>A</i> <sup>v</sup>  | 3.108 (9) |
| Cs1—O14 <i>B</i> <sup>iii</sup> | 3.183 (9) | Cs2—O14 <i>B</i> <sup>v</sup>  | 3.130 (9) |

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $-x + 1, y, -z + \frac{1}{2}$ ; (iii)  $-x + 1, y + 1, -z + \frac{1}{2}$ ; (iv)  $-x, y, -z + \frac{1}{2}$ ; (v)  $-x, y + 1, -z + \frac{1}{2}$ .

**Table 2**

Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$                                   | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|--|-------|--------------|--------------|----------------|
| O14 <i>B</i> —H14 <i>B</i> $\cdots$ O14 <i>A</i> | 1.21  | 1.25         | 2.462 (10)   | 180            |

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *WinGX* (Farrugia, 2012); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

The author acknowledges financial support from the Science and Engineering Faculty and the University Library, Queensland University of Technology.

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM2798).

## References

- Abdelmoty, I., Bucholz, V., Di, L., Guo, C., Kowitz, K., Enkelmann, V., Wegner, G. & Foxman, B. M. (2005). *Cryst. Growth Des.* **5**, 2210–2217.
- Agilent (2013). *CrysAlis PRO*. Agilent Technologies Ltd, Yarnton, England.
- Chowdhury, M. & Kariuki, B. M. (2006). *Cryst. Growth Des.* **6**, 774–780.
- Crowther, D., Chowdhury, M. & Kariuki, B. M. (2008). *J. Mol. Struct.* **872**, 64–71.

- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Kariuki, B. M., Valim, J. B., Jones, W. & King, J. (1994). *Acta Cryst. C* **50**, 1665–1667.
- Kariuki, B. M., Valim, J. B., Jones, W. & King, J. (1995). *Acta Cryst. C* **51**, 1051–1053.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Smith, G. & Wermuth, U. D. (2009). *Acta Cryst. E* **65**, m1048.
- Smith, G. & Wermuth, U. D. (2011). *Acta Cryst. E* **67**, m1594–m1595.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wierda, D. A., Feng, T. L. & Barron, A. R. (1989). *Acta Cryst. C* **45**, 338–339.

## supplementary materials

*Acta Cryst.* (2014). E70, m43–m44 [doi:10.1107/S1600536814000804]

## Poly[ $\mu_5$ -{hydrogen bis[(*E*)-cinnamato]}-caesium]

Graham Smith

### 1. Comment

The crystal structure of *trans*-cinnamic acid was reported by Wierda *et al.* (1989) and Abdelmoty *et al.* (2005). The alkali metal salts of *trans*-cinnamic acid are unknown in the crystallographic literature although a limited number of examples of salts of ring-substituted cinnamates have been reported, *e.g.* the sodium salts of 2-nitrocinnamate [a dihydrate (Smith & Wermuth, 2009)], of 2-chlorocinnamate [a dihydrate (Kariuki *et al.*, 1995)], of 3-chlorocinnamate [anhydrous (Crowther *et al.*, 2008)], of 4-chlorocinnamate [a dihydrate (Kariuki *et al.*, 1994)]; potassium salts of 3-chloro- and 3-bromocinnamate [both anhydrous (Crowther *et al.*, 2008)]; and a rubidium salt of 2-nitrocinnamate [a monohydrate (Smith & Wermuth, 2011)].

The reaction of *trans*-cinnamic acid with caesium hydroxide in aqueous ethanol afforded crystals of the title complex, [Cs(C<sub>9</sub>H<sub>7</sub>O<sub>2</sub>)(C<sub>9</sub>H<sub>8</sub>O<sub>2</sub>)]<sub>n</sub>, (I), the structure of which is reported herein.

In the structure of (I) the asymmetric unit (Fig. 1) comprises two independent irregular CsO<sub>8</sub> coordination polyhedra [Cs1—O, 3.060 (8)–3.183 (9) Å; Cs2—O, 3.063 (9)–3.377 (9) Å; Table 1], in which the Cs<sup>+</sup> ions lie on a twofold rotation axis and are linked by four bridging carboxyl O-donors from the two *trans*-cinnamate ligand species. These two ligand species are inter-linked through a delocalized H atom on an approximately central intermediate site within a short O14A···H14B···O14B hydrogen bond [2.462 (10) Å] (Table 2). Although this phenomenon involving coordinating dimeric carboxylate species is not known among the alkali metal substituted-cinnamate structures, it is found in both ammonium hydrogen bis(3-chlorocinnamate) and ammonium hydrogen bis(3-bromocinnamate) (Chowdhury & Kariuki, 2006), with the O···H···O values [2.554 (6) Å for the 3-Cl-analogue and 2.466 (5) Å for the 3-Br-analogue] similar to that in the structure of (I). In this complex, the two Cs<sup>+</sup> ions are quadruply bridged giving a Cs1···Cs2 separation of 3.9318 (3) Å and generate an overall two-dimensional coordination polymer lying parallel to (001) (Figs. 2, 3). No inter-ring  $\pi$ – $\pi$  interactions are present in the structure [minimum ring centroid separation = 4.826 (8) Å].

The two linked cinnamate species in the title complex are close to coplanar [inter-ring dihedral angle = 3.9 (6)°], with the side chain carboxyl group of the *A* ligand component slightly rotated out of the plane [torsion angle C11A—C12A—C13A—O13A = 169.0 (13)°] compared to that of the *B* ligand component [torsion angle C11B—C12B—C13B—O14B = -179.2 (11)°]. With the analogous ammonium hydrogen salts of the 3-chloro- and 3-bromocinnamates (Chowdhury & Kariuki, 2006), the two cinnamate components are related either by crystallographic inversion symmetry (3-Cl) with the two benzene rings essentially planar, or by twofold rotational symmetry (3-Br) with the two rings significantly rotated out of the least-squares plane [inter-ring dihedral angle = 29.8 (2)°].

### 2. Experimental

The title compound was synthesized by heating together for 10 minutes, 148 mg (1.0 mmol) of *trans*-cinnamic acid and 75 mg (0.5 mmol) of CsOH in 15 ml of an 1:9 (vol/vol) ethanol–water mixture. Partial room temperature evaporation of the solution gave colourless elongated crystals of the title complex from which a specimen was cleaved for the X-ray


analysis. These crystals were invariably twinned, a feature identified in the later structure solution and refinement routines.

### 3. Refinement

Hydrogen atoms were placed in calculated positions [ $C-H = 0.95 \text{ \AA}$ ] and allowed to ride in the refinement, with  $U_{iso}(H) = 1.2U_{eq}(C)$ . The carboxylic acid H-atom was found to be delocalized in a site approximating to midway between two carboxyl O-atoms of the dimeric acid-anion unit and was subsequently allowed to ride at that site, with  $U_{iso}(H) = 1.5U_{eq}(O)$ . The presence of a non-merohedral twin was identified using TwinRotMat within *PLATON* (Spek, 2009) (twin law:  $\bar{1} 0 0, 0 \bar{1} 0, 1.5 0 1$ ) reducing the conventional *R*-factor from 0.23 to 0.072, with a final BASF factor (HKL 5 format) of 0.4836. Maximum and minimum residual electron densities were  $1.26 \text{ e \AA}^{-3}$  ( $1.00 \text{ \AA}$  from Cs1) and  $-2.19 \text{ e \AA}^{-3}$  ( $1.94 \text{ \AA}$  from H14B), respectively.


### Computing details

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *WinGX* (Farrugia, 2012); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON* (Spek, 2009).




**Figure 1**

The atom-numbering scheme and the molecular configuration of the two ligands and the two  $CsO_8$  coordination polyhedra of the title complex, with non-H atoms drawn with displacement ellipsoids at the 40% probability level. The two  $Cs^+$  cations lie on twofold rotation axes. The  $O14A \cdots O14B$  hydrogen bond with the delocalized H atom (H14B) is shown as a dashed link. [For symmetry codes: see Table 1].

**Figure 2**

A view of the partially expanded polymeric extension of the structure viewed along the approximate *a*-cell direction. C-bound H atoms are omitted. A and B denote the two different ligand components.

**Figure 3**

The packing of the layered structure of compound (I) viewed along *b*.

### Poly[ $\mu_5$ -{hydrogen bis[(*E*)-cinnamato}]-caesium]

#### Crystal data

[Cs(C<sub>9</sub>H<sub>7</sub>O<sub>2</sub>)(C<sub>9</sub>H<sub>8</sub>O<sub>2</sub>)]

$M_r = 428.21$

Monoclinic, *P*2/*c*

Hall symbol: -P 2yc

$a = 7.8608 (6) \text{ \AA}$

$b = 5.6985 (7) \text{ \AA}$

$c = 38.817 (3) \text{ \AA}$

$\beta = 98.733 (6)^\circ$

$V = 1718.6 (3) \text{ \AA}^3$   
 $Z = 4$   
 $F(000) = 840$   
 $D_x = 1.655 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 1674 reflections

$\theta = 3.6\text{--}28.2^\circ$   
 $\mu = 2.17 \text{ mm}^{-1}$   
 $T = 200 \text{ K}$   
Plate, colourless  
 $0.35 \times 0.35 \times 0.06 \text{ mm}$

#### Data collection

Oxford Diffraction Gemini-S CCD-detector  
diffractometer  
Radiation source: Enhance (Mo) X-ray source  
Graphite monochromator  
Detector resolution:  $16.077 \text{ pixels mm}^{-1}$   
 $\omega$  scans  
Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent, 2013)  
 $T_{\min} = 0.711$ ,  $T_{\max} = 0.980$

6675 measured reflections  
3353 independent reflections  
2552 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$   
 $\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 3.2^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -7 \rightarrow 7$   
 $l = -11 \rightarrow 47$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.071$   
 $wR(F^2) = 0.144$   
 $S = 1.19$   
3353 reflections  
210 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.020P)^2 + 18.34P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.26 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -2.19 \text{ e \AA}^{-3}$

#### Special details

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

|      | <i>x</i>     | <i>y</i>     | <i>z</i>     | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|--------------|--------------|--------------|----------------------------------|
| Cs1  | 0.50000      | 0.6438 (2)   | 0.25000      | 0.0261 (3)                       |
| Cs2  | 0.00000      | 0.6257 (2)   | 0.25000      | 0.0300 (3)                       |
| O13A | 0.2828 (13)  | −0.1322 (15) | 0.3028 (2)   | 0.037 (3)                        |
| O13B | 0.2180 (12)  | 0.3903 (14)  | 0.20079 (19) | 0.030 (3)                        |
| O14A | 0.3036 (12)  | 0.2355 (14)  | 0.28374 (18) | 0.029 (3)                        |
| O14B | 0.2197 (13)  | 0.0396 (14)  | 0.22718 (18) | 0.033 (3)                        |
| C1A  | 0.5037 (15)  | 0.4316 (19)  | 0.3905 (3)   | 0.025 (3)                        |
| C1B  | 0.0000 (15)  | 0.079 (2)    | 0.1031 (3)   | 0.025 (3)                        |
| C2A  | 0.5963 (19)  | 0.636 (2)    | 0.3964 (3)   | 0.035 (4)                        |
| C2B  | 0.0132 (16)  | 0.208 (2)    | 0.0728 (3)   | 0.028 (3)                        |
| C3A  | 0.6620 (19)  | 0.707 (2)    | 0.4301 (4)   | 0.042 (5)                        |
| C3B  | −0.0664 (16) | 0.128 (3)    | 0.0403 (3)   | 0.040 (4)                        |



|      |              |            |            |           |
|------|--------------|------------|------------|-----------|
| C4A  | 0.6346 (18)  | 0.574 (3)  | 0.4585 (3) | 0.042 (5) |
| C4B  | −0.1524 (18) | −0.082 (2) | 0.0373 (3) | 0.041 (4) |
| C5A  | 0.5426 (19)  | 0.373 (3)  | 0.4529 (3) | 0.039 (4) |
| C5B  | −0.1628 (16) | −0.210 (2) | 0.0668 (3) | 0.035 (4) |
| C6A  | 0.4764 (15)  | 0.299 (2)  | 0.4191 (3) | 0.029 (4) |
| C6B  | −0.0870 (15) | −0.133 (2) | 0.0997 (3) | 0.031 (4) |
| C11A | 0.4356 (17)  | 0.359 (2)  | 0.3541 (3) | 0.032 (4) |
| C11B | 0.0831 (14)  | 0.173 (2)  | 0.1369 (3) | 0.025 (3) |
| C12A | 0.3722 (13)  | 0.155 (2)  | 0.3444 (3) | 0.023 (3) |
| C12B | 0.1178 (15)  | 0.054 (2)  | 0.1672 (3) | 0.026 (3) |
| C13A | 0.3148 (16)  | 0.076 (2)  | 0.3078 (3) | 0.028 (4) |
| C13B | 0.1901 (15)  | 0.176 (2)  | 0.1998 (3) | 0.027 (4) |
| H2A  | 0.61560      | 0.73090    | 0.37720    | 0.0410*   |
| H2B  | 0.07640      | 0.35110    | 0.07440    | 0.0340*   |
| H3A  | 0.72650      | 0.84820    | 0.43360    | 0.0510*   |
| H3B  | −0.06090     | 0.21990    | 0.02000    | 0.0480*   |
| H4A  | 0.67930      | 0.62310    | 0.48150    | 0.0500*   |
| H4B  | −0.20410     | −0.13800   | 0.01520    | 0.0490*   |
| H5A  | 0.52280      | 0.28000    | 0.47230    | 0.0470*   |
| H5B  | −0.22320     | −0.35490   | 0.06480    | 0.0410*   |
| H6A  | 0.41240      | 0.15700    | 0.41580    | 0.0350*   |
| H6B  | −0.09500     | −0.22560   | 0.11970    | 0.0370*   |
| H11A | 0.43910      | 0.47280    | 0.33640    | 0.0380*   |
| H11B | 0.11540      | 0.33430    | 0.13740    | 0.0290*   |
| H12A | 0.36110      | 0.04540    | 0.36230    | 0.0280*   |
| H12B | 0.09570      | −0.10990   | 0.16760    | 0.0320*   |
| H14B | 0.26080      | 0.13620    | 0.25500    | 0.0500*   |

Atomic displacement parameters ( $\text{\AA}^2$ )

|      | $U^{11}$   | $U^{22}$   | $U^{33}$   | $U^{12}$   | $U^{13}$   | $U^{23}$   |
|------|------------|------------|------------|------------|------------|------------|
| Cs1  | 0.0221 (5) | 0.0212 (5) | 0.0350 (5) | 0.0000     | 0.0041 (5) | 0.0000     |
| Cs2  | 0.0232 (5) | 0.0238 (6) | 0.0438 (6) | 0.0000     | 0.0075 (6) | 0.0000     |
| O13A | 0.056 (5)  | 0.027 (5)  | 0.028 (4)  | −0.001 (6) | 0.003 (4)  | −0.006 (4) |
| O13B | 0.047 (5)  | 0.026 (4)  | 0.016 (4)  | −0.004 (5) | 0.000 (4)  | 0.000 (3)  |
| O14A | 0.046 (6)  | 0.033 (5)  | 0.006 (3)  | −0.003 (4) | −0.006 (3) | −0.005 (3) |
| O14B | 0.055 (5)  | 0.025 (4)  | 0.013 (4)  | −0.004 (5) | −0.015 (4) | −0.005 (3) |
| C1A  | 0.022 (6)  | 0.022 (6)  | 0.030 (6)  | 0.003 (5)  | 0.006 (5)  | −0.009 (5) |
| C1B  | 0.020 (5)  | 0.034 (6)  | 0.022 (5)  | 0.004 (6)  | 0.005 (5)  | −0.007 (5) |
| C2A  | 0.038 (7)  | 0.027 (6)  | 0.039 (6)  | −0.004 (7) | 0.007 (6)  | 0.003 (6)  |
| C2B  | 0.027 (6)  | 0.028 (6)  | 0.027 (6)  | 0.007 (6)  | 0.000 (5)  | −0.003 (5) |
| C3A  | 0.044 (9)  | 0.021 (7)  | 0.059 (9)  | 0.012 (6)  | −0.002 (7) | −0.024 (6) |
| C3B  | 0.043 (8)  | 0.050 (8)  | 0.027 (6)  | 0.026 (8)  | 0.005 (5)  | 0.000 (7)  |
| C4A  | 0.031 (7)  | 0.059 (10) | 0.033 (7)  | 0.006 (8)  | −0.001 (6) | −0.022 (7) |
| C4B  | 0.037 (7)  | 0.053 (8)  | 0.030 (7)  | −0.011 (8) | −0.003 (6) | −0.012 (6) |
| C5A  | 0.035 (7)  | 0.048 (7)  | 0.035 (6)  | −0.003 (8) | 0.005 (6)  | 0.012 (7)  |
| C5B  | 0.029 (7)  | 0.021 (6)  | 0.054 (8)  | −0.005 (6) | 0.006 (6)  | −0.004 (6) |
| C6A  | 0.031 (7)  | 0.027 (7)  | 0.030 (6)  | −0.007 (6) | 0.005 (5)  | 0.004 (5)  |
| C6B  | 0.032 (7)  | 0.027 (6)  | 0.032 (6)  | −0.002 (6) | 0.002 (5)  | −0.002 (6) |
| C11A | 0.035 (7)  | 0.029 (6)  | 0.032 (6)  | −0.002 (7) | 0.006 (6)  | 0.002 (6)  |

|      |           |           |           |            |            |            |
|------|-----------|-----------|-----------|------------|------------|------------|
| C11B | 0.030 (6) | 0.021 (6) | 0.021 (5) | −0.004 (5) | −0.001 (4) | 0.001 (5)  |
| C12A | 0.036 (6) | 0.017 (6) | 0.016 (5) | −0.008 (5) | 0.002 (4)  | −0.008 (5) |
| C12B | 0.038 (7) | 0.013 (5) | 0.027 (6) | −0.009 (5) | 0.001 (5)  | 0.006 (5)  |
| C13A | 0.021 (6) | 0.036 (8) | 0.027 (6) | −0.002 (6) | 0.003 (5)  | 0.002 (5)  |
| C13B | 0.023 (6) | 0.043 (8) | 0.015 (5) | 0.000 (6)  | −0.001 (5) | −0.010 (6) |

*Geometric parameters (Å, °)*

|  |             |   |            |
|--|-------------|---|------------|
| Cs1—O13B                                 | 3.060 (8)   | C2A—C3A                                   | 1.392 (19) |
| Cs1—O14A                                 | 3.182 (8)   | C2B—C3B                                   | 1.397 (17) |
| Cs1—O13A <sup>i</sup>                    | 3.132 (9)   | C3A—C4A                                   | 1.38 (2)   |
| Cs1—O14B <sup>i</sup>                    | 3.183 (9)   | C3B—C4B                                   | 1.37 (2)   |
| Cs1—O13B <sup>ii</sup>                   | 3.060 (8)   | C4A—C5A                                   | 1.35 (2)   |
| Cs1—O14A <sup>ii</sup>                   | 3.182 (8)   | C4B—C5B                                   | 1.371 (16) |
| Cs1—O13A <sup>iii</sup>                  | 3.132 (9)   | C5A—C6A                                   | 1.401 (17) |
| Cs1—O14B <sup>iii</sup>                  | 3.183 (9)   | C5B—C6B                                   | 1.396 (16) |
| Cs2—O13B                                 | 3.063 (8)   | C11A—C12A                                 | 1.298 (16) |
| Cs2—O14A                                 | 3.377 (9)   | C11B—C12B                                 | 1.349 (16) |
| Cs2—O13A <sup>i</sup>                    | 3.108 (9)   | C12A—C13A                                 | 1.493 (16) |
| Cs2—O14B <sup>i</sup>                    | 3.130 (9)   | C12B—C13B                                 | 1.480 (16) |
| Cs2—O13B <sup>iv</sup>                   | 3.063 (8)   | C2A—H2A                                   | 0.9500     |
| Cs2—O14A <sup>iv</sup>                   | 3.377 (9)   | C2B—H2B                                   | 0.9500     |
| Cs2—O13A <sup>v</sup>                    | 3.108 (9)   | C3A—H3A                                   | 0.9500     |
| Cs2—O14B <sup>v</sup>                    | 3.130 (9)   | C3B—H3B                                   | 0.9500     |
| O13A—C13A                                | 1.222 (14)  | C4A—H4A                                   | 0.9500     |
| O13B—C13B                                | 1.240 (14)  | C4B—H4B                                   | 0.9500     |
| O14A—C13A                                | 1.297 (14)  | C5A—H5A                                   | 0.9500     |
| O14B—C13B                                | 1.309 (14)  | C5B—H5B                                   | 0.9500     |
| O14B—H14B                                | 1.2100      | C6A—H6A                                   | 0.9500     |
| C1A—C11A                                 | 1.492 (16)  | C6B—H6B                                   | 0.9500     |
| C1A—C2A                                  | 1.374 (17)  | C11A—H11A                                 | 0.9500     |
| C1A—C6A                                  | 1.386 (16)  | C11B—H11B                                 | 0.9500     |
| C1B—C11B                                 | 1.475 (16)  | C12A—H12A                                 | 0.9500     |
| C1B—C6B                                  | 1.385 (16)  | C12B—H12B                                 | 0.9500     |
| C1B—C2B                                  | 1.404 (16)  |   |            |
| O13B—Cs1—O14A                            | 64.0 (2)    | Cs2—O14A—C13A                             | 135.0 (8)  |
| O13A <sup>i</sup> —Cs1—O13B              | 100.7 (2)   | Cs1 <sup>vi</sup> —O14B—C13B              | 132.6 (7)  |
| O13B—Cs1—O14B <sup>i</sup>               | 75.9 (2)    | Cs2 <sup>vi</sup> —O14B—C13B              | 129.8 (7)  |
| O13B—Cs1—O13B <sup>ii</sup>              | 123.7 (2)   | Cs1 <sup>vi</sup> —O14B—Cs2 <sup>vi</sup> | 77.04 (18) |
| O13B—Cs1—O14A <sup>ii</sup>              | 75.4 (2)    | Cs1—O14A—H14B                             | 93.00      |
| O13A <sup>iii</sup> —Cs1—O13B            | 101.5 (2)   | Cs2—O14A—H14B                             | 83.00      |
| O13B—Cs1—O14B <sup>iii</sup>             | 155.98 (19) | Cs2 <sup>vi</sup> —O14B—H14B              | 100.00     |
| O13A <sup>i</sup> —Cs1—O14A              | 71.5 (2)    | Cs1 <sup>vi</sup> —O14B—H14B              | 90.00      |
| O14A—Cs1—O14B <sup>i</sup>               | 105.9 (2)   | C13B—O14B—H14B                            | 116.00     |
| O13B <sup>ii</sup> —Cs1—O14A             | 75.4 (2)    | C6A—C1A—C11A                              | 122.0 (10) |
| O14A—Cs1—O14A <sup>ii</sup>              | 86.0 (2)    | C2A—C1A—C6A                               | 118.1 (11) |
| O13A <sup>iii</sup> —Cs1—O14A            | 156.1 (2)   | C2A—C1A—C11A                              | 119.9 (10) |
| O14A—Cs1—O14B <sup>iii</sup>             | 139.65 (18) | C2B—C1B—C6B                               | 118.4 (11) |
| O13A <sup>i</sup> —Cs1—O14B <sup>i</sup> | 58.0 (2)    | C2B—C1B—C11B                              | 118.4 (10) |

|  |             |                |            |
|--|-------------|----------------|------------|
| O13A <sup>i</sup> —Cs1—O13B <sup>ii</sup>    | 101.5 (2)   | C6B—C1B—C11B   | 123.2 (10) |
| O13A <sup>i</sup> —Cs1—O14A <sup>ii</sup>    | 156.1 (2)   | C1A—C2A—C3A    | 121.0 (11) |
| O13A <sup>i</sup> —Cs1—O13A <sup>iii</sup>   | 131.9 (2)   | C1B—C2B—C3B    | 120.4 (12) |
| O13A <sup>i</sup> —Cs1—O14B <sup>iii</sup>   | 87.3 (2)    | C2A—C3A—C4A    | 120.7 (12) |
| O13B <sup>ii</sup> —Cs1—O14B <sup>i</sup>    | 155.98 (19) | C2B—C3B—C4B    | 120.6 (12) |
| O14A <sup>ii</sup> —Cs1—O14B <sup>i</sup>    | 139.65 (18) | C3A—C4A—C5A    | 118.7 (12) |
| O13A <sup>iii</sup> —Cs1—O14B <sup>i</sup>   | 87.3 (2)    | C3B—C4B—C5B    | 119.0 (11) |
| O14B <sup>i</sup> —Cs1—O14B <sup>iii</sup>   | 89.8 (2)    | C4A—C5A—C6A    | 121.2 (12) |
| O13B <sup>ii</sup> —Cs1—O14A <sup>ii</sup>   | 64.0 (2)    | C4B—C5B—C6B    | 121.7 (11) |
| O13A <sup>iii</sup> —Cs1—O13B <sup>ii</sup>  | 100.7 (2)   | C1A—C6A—C5A    | 120.4 (11) |
| O13B <sup>ii</sup> —Cs1—O14B <sup>iii</sup>  | 75.9 (2)    | C1B—C6B—C5B    | 119.8 (11) |
| O13A <sup>iii</sup> —Cs1—O14A <sup>ii</sup>  | 71.5 (2)    | C1A—C11A—C12A  | 126.2 (11) |
| O14A <sup>ii</sup> —Cs1—O14B <sup>iii</sup>  | 105.9 (2)   | C1B—C11B—C12B  | 126.6 (11) |
| O13A <sup>iii</sup> —Cs1—O14B <sup>iii</sup> | 58.0 (2)    | C11A—C12A—C13A | 126.4 (11) |
| O13B—Cs2—O14A                                | 61.58 (19)  | C11B—C12B—C13B | 120.7 (10) |
| O13A <sup>i</sup> —Cs2—O13B                  | 101.2 (2)   | O13A—C13A—C12A | 118.0 (10) |
| O13B—Cs2—O14B <sup>i</sup>                   | 76.6 (2)    | O13A—C13A—O14A | 125.2 (11) |
| O13B—Cs2—O13B <sup>iv</sup>                  | 128.1 (2)   | O14A—C13A—C12A | 116.9 (10) |
| O13B—Cs2—O14A <sup>iv</sup>                  | 84.2 (2)    | O14B—C13B—C12B | 114.4 (10) |
| O13A <sup>v</sup> —Cs2—O13B                  | 101.3 (2)   | O13B—C13B—O14B | 123.4 (10) |
| O13B—Cs2—O14B <sup>v</sup>                   | 153.1 (2)   | O13B—C13B—C12B | 122.2 (10) |
| O13A <sup>i</sup> —Cs2—O14A                  | 69.2 (2)    | C1A—C2A—H2A    | 120.00     |
| O14A—Cs2—O14B <sup>i</sup>                   | 102.6 (2)   | C3A—C2A—H2A    | 119.00     |
| O13B <sup>iv</sup> —Cs2—O14A                 | 84.2 (2)    | C1B—C2B—H2B    | 120.00     |
| O14A—Cs2—O14A <sup>iv</sup>                  | 97.7 (2)    | C3B—C2B—H2B    | 120.00     |
| O13A <sup>v</sup> —Cs2—O14A                  | 160.1 (2)   | C2A—C3A—H3A    | 120.00     |
| O14A—Cs2—O14B <sup>v</sup>                   | 140.74 (18) | C4A—C3A—H3A    | 120.00     |
| O13A <sup>i</sup> —Cs2—O14B <sup>i</sup>     | 58.8 (2)    | C2B—C3B—H3B    | 120.00     |
| O13A <sup>i</sup> —Cs2—O13B <sup>iv</sup>    | 101.3 (2)   | C4B—C3B—H3B    | 120.00     |
| O13A <sup>i</sup> —Cs2—O14A <sup>iv</sup>    | 160.1 (2)   | C3A—C4A—H4A    | 121.00     |
| O13A <sup>i</sup> —Cs2—O13A <sup>v</sup>     | 127.3 (2)   | C5A—C4A—H4A    | 121.00     |
| O13A <sup>i</sup> —Cs2—O14B <sup>v</sup>     | 81.3 (2)    | C3B—C4B—H4B    | 121.00     |
| O13B <sup>iv</sup> —Cs2—O14B <sup>i</sup>    | 153.1 (2)   | C5B—C4B—H4B    | 120.00     |
| O14A <sup>iv</sup> —Cs2—O14B <sup>i</sup>    | 140.74 (18) | C4A—C5A—H5A    | 119.00     |
| O13A <sup>v</sup> —Cs2—O14B <sup>i</sup>     | 81.3 (2)    | C6A—C5A—H5A    | 119.00     |
| O14B <sup>i</sup> —Cs2—O14B <sup>v</sup>     | 82.2 (2)    | C4B—C5B—H5B    | 119.00     |
| O13B <sup>iv</sup> —Cs2—O14A <sup>iv</sup>   | 61.58 (19)  | C6B—C5B—H5B    | 119.00     |
| O13A <sup>v</sup> —Cs2—O13B <sup>iv</sup>    | 101.2 (2)   | C1A—C6A—H6A    | 120.00     |
| O13B <sup>iv</sup> —Cs2—O14B <sup>v</sup>    | 76.6 (2)    | C5A—C6A—H6A    | 120.00     |
| O13A <sup>v</sup> —Cs2—O14A <sup>iv</sup>    | 69.2 (2)    | C1B—C6B—H6B    | 120.00     |
| O14A <sup>iv</sup> —Cs2—O14B <sup>v</sup>    | 102.6 (2)   | C5B—C6B—H6B    | 120.00     |
| O13A <sup>v</sup> —Cs2—O14B <sup>v</sup>     | 58.8 (2)    | C1A—C11A—H11A  | 117.00     |
| Cs1 <sup>vi</sup> —O13A—C13A                 | 112.3 (8)   | C12A—C11A—H11A | 117.00     |
| Cs2 <sup>vi</sup> —O13A—C13A                 | 130.2 (8)   | C1B—C11B—H11B  | 117.00     |
| Cs1 <sup>vi</sup> —O13A—Cs2 <sup>vi</sup>    | 78.11 (18)  | C12B—C11B—H11B | 117.00     |
| Cs1—O13B—Cs2                                 | 79.91 (18)  | C11A—C12A—H12A | 117.00     |
| Cs1—O13B—C13B                                | 126.3 (7)   | C13A—C12A—H12A | 117.00     |
| Cs2—O13B—C13B                                | 109.7 (7)   | C11B—C12B—H12B | 120.00     |
| Cs1—O14A—Cs2                                 | 73.59 (16)  | C13B—C12B—H12B | 120.00     |

|                                    |              |                                   |              |
|------------------------------------|--------------|-----------------------------------|--------------|
| Cs1—O14A—C13A                      | 145.6 (8)    |                                   |              |
| O14A—Cs1—O13B—Cs2                  | 65.6 (2)     | O13B <sup>iv</sup> —Cs2—O14A—Cs1  | −160.72 (18) |
| O14A—Cs1—O13B—C13B                 | −41.6 (9)    | O13B <sup>iv</sup> —Cs2—O14A—C13A | −3.5 (9)     |
| O13A <sup>i</sup> —Cs1—O13B—Cs2    | 2.5 (2)      | O14A <sup>iv</sup> —Cs2—O14A—Cs1  | 139.03 (16)  |
| O13A <sup>i</sup> —Cs1—O13B—C13B   | −104.7 (9)   | O14A <sup>iv</sup> —Cs2—O14A—C13A | −63.8 (10)   |
| O14B <sup>i</sup> —Cs1—O13B—Cs2    | −50.38 (19)  | O14B <sup>v</sup> —Cs2—O14A—Cs1   | −100.2 (3)   |
| O14B <sup>i</sup> —Cs1—O13B—C13B   | −157.5 (9)   | O14B <sup>v</sup> —Cs2—O14A—C13A  | 57.0 (11)    |
| O13B <sup>ii</sup> —Cs1—O13B—Cs2   | 114.2 (2)    | O13B—Cs2—O13A <sup>i</sup> —Cs1   | 2.5 (2)      |
| O13B <sup>ii</sup> —Cs1—O13B—C13B  | 7.0 (10)     | O14A—Cs2—O13A <sup>i</sup> —Cs1   | 56.01 (18)   |
| O14A <sup>ii</sup> —Cs1—O13B—Cs2   | 158.3 (2)    | O13B—Cs2—O14B <sup>i</sup> —Cs1   | −49.32 (18)  |
| O14A <sup>ii</sup> —Cs1—O13B—C13B  | 51.2 (9)     | O14A—Cs2—O14B <sup>i</sup> —Cs1   | 7.01 (18)    |
| O13A <sup>iii</sup> —Cs1—O13B—Cs2  | −134.6 (2)   | Cs1 <sup>vi</sup> —O13A—C13A—O14A | 56.2 (15)    |
| O13A <sup>iii</sup> —Cs1—O13B—C13B | 118.3 (9)    | Cs1 <sup>vi</sup> —O13A—C13A—C12A | −123.5 (9)   |
| O14B <sup>iii</sup> —Cs1—O13B—Cs2  | −105.2 (5)   | Cs2 <sup>vi</sup> —O13A—C13A—O14A | −37.0 (18)   |
| O14B <sup>iii</sup> —Cs1—O13B—C13B | 147.6 (9)    | Cs2 <sup>vi</sup> —O13A—C13A—C12A | 143.3 (8)    |
| O13B—Cs1—O14A—Cs2                  | −57.9 (2)    | Cs1—O13B—C13B—O14B                | 31.2 (16)    |
| O13B—Cs1—O14A—C13A                 | 151.0 (13)   | Cs1—O13B—C13B—C12B                | −149.6 (8)   |
| O13A <sup>i</sup> —Cs1—O14A—Cs2    | 54.52 (19)   | Cs2—O13B—C13B—O14B                | −60.7 (13)   |
| O13A <sup>i</sup> —Cs1—O14A—C13A   | −96.5 (12)   | Cs2—O13B—C13B—C12B                | 118.6 (10)   |
| O14B <sup>i</sup> —Cs1—O14A—Cs2    | 7.10 (18)    | Cs1—O14A—C13A—O13A                | −125.8 (12)  |
| O14B <sup>i</sup> —Cs1—O14A—C13A   | −143.9 (12)  | Cs1—O14A—C13A—C12A                | 54.0 (17)    |
| O13B <sup>ii</sup> —Cs1—O14A—Cs2   | 162.23 (19)  | Cs2—O14A—C13A—O13A                | 95.3 (14)    |
| O13B <sup>ii</sup> —Cs1—O14A—C13A  | 11.2 (12)    | Cs2—O14A—C13A—C12A                | −85.0 (13)   |
| O14A <sup>ii</sup> —Cs1—O14A—Cs2   | −133.66 (17) | Cs1 <sup>vi</sup> —O14B—C13B—O13B | −109.4 (12)  |
| O14A <sup>ii</sup> —Cs1—O14A—C13A  | 75.3 (12)    | Cs1 <sup>vi</sup> —O14B—C13B—C12B | 71.3 (13)    |
| O13A <sup>iii</sup> —Cs1—O14A—Cs2  | −114.2 (5)   | Cs2 <sup>vi</sup> —O14B—C13B—O13B | 138.9 (10)   |
| O13A <sup>iii</sup> —Cs1—O14A—C13A | 94.7 (13)    | Cs2 <sup>vi</sup> —O14B—C13B—C12B | −40.4 (14)   |
| O14B <sup>iii</sup> —Cs1—O14A—Cs2  | 116.3 (3)    | C6A—C1A—C2A—C3A                   | 0.5 (19)     |
| O14B <sup>iii</sup> —Cs1—O14A—C13A | −34.7 (14)   | C11A—C1A—C2A—C3A                  | −179.7 (12)  |
| O13B—Cs1—O13A <sup>i</sup> —Cs2    | −2.5 (2)     | C2A—C1A—C6A—C5A                   | −0.2 (18)    |
| O14A—Cs1—O13A <sup>i</sup> —Cs2    | −60.15 (19)  | C11A—C1A—C6A—C5A                  | 179.9 (12)   |
| O13B—Cs1—O14B <sup>i</sup> —Cs2    | 49.60 (18)   | C2A—C1A—C11A—C12A                 | 167.7 (13)   |
| O14A—Cs1—O14B <sup>i</sup> —Cs2    | −7.55 (19)   | C6A—C1A—C11A—C12A                 | −13 (2)      |
| O14A—Cs2—O13B—Cs1                  | −61.2 (2)    | C6B—C1B—C2B—C3B                   | 2.8 (18)     |
| O14A—Cs2—O13B—C13B                 | 63.9 (7)     | C11B—C1B—C2B—C3B                  | −178.7 (11)  |
| O13A <sup>i</sup> —Cs2—O13B—Cs1    | −2.5 (2)     | C2B—C1B—C6B—C5B                   | −1.7 (18)    |
| O13A <sup>i</sup> —Cs2—O13B—C13B   | 122.6 (7)    | C11B—C1B—C6B—C5B                  | 179.8 (11)   |
| O14B <sup>i</sup> —Cs2—O13B—Cs1    | 51.35 (19)   | C2B—C1B—C11B—C12B                 | −164.0 (12)  |
| O14B <sup>i</sup> —Cs2—O13B—C13B   | 176.5 (8)    | C6B—C1B—C11B—C12B                 | 14.4 (19)    |
| O13B <sup>iv</sup> —Cs2—O13B—Cs1   | −116.5 (2)   | C1A—C2A—C3A—C4A                   | 0 (2)        |
| O13B <sup>iv</sup> —Cs2—O13B—C13B  | 8.6 (8)      | C1B—C2B—C3B—C4B                   | −3 (2)       |
| O14A <sup>iv</sup> —Cs2—O13B—Cs1   | −163.16 (19) | C2A—C3A—C4A—C5A                   | 0 (2)        |
| O14A <sup>iv</sup> —Cs2—O13B—C13B  | −38.1 (7)    | C2B—C3B—C4B—C5B                   | 2 (2)        |
| O13A <sup>v</sup> —Cs2—O13B—Cs1    | 129.5 (2)    | C3A—C4A—C5A—C6A                   | 0 (2)        |
| O13A <sup>v</sup> —Cs2—O13B—C13B   | −105.4 (7)   | C3B—C4B—C5B—C6B                   | −1 (2)       |
| O14B <sup>v</sup> —Cs2—O13B—Cs1    | 90.3 (5)     | C4A—C5A—C6A—C1A                   | 0 (2)        |
| O14B <sup>v</sup> —Cs2—O13B—C13B   | −144.6 (7)   | C4B—C5B—C6B—C1B                   | 0.7 (19)     |
| O13B—Cs2—O14A—Cs1                  | 59.9 (2)     | C1A—C11A—C12A—C13A                | −175.6 (12)  |

|                                  |             |                     |             |
|----------------------------------|-------------|---------------------|-------------|
| O13B—Cs2—O14A—C13A               | −142.9 (10) | C1B—C11B—C12B—C13B  | −175.3 (11) |
| O13A <sup>i</sup> —Cs2—O14A—Cs1  | −56.3 (2)   | C11A—C12A—C13A—O13A | 169.0 (13)  |
| O13A <sup>i</sup> —Cs2—O14A—C13A | 100.9 (10)  | C11A—C12A—C13A—O14A | −10.7 (18)  |
| O14B <sup>i</sup> —Cs2—O14A—Cs1  | −7.12 (18)  | C11B—C12B—C13B—O13B | 1.6 (18)    |
| O14B <sup>i</sup> —Cs2—O14A—C13A | 150.1 (9)   | C11B—C12B—C13B—O14B | −179.2 (11) |

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $-x+1, y, -z+1/2$ ; (iii)  $-x+1, y+1, -z+1/2$ ; (iv)  $-x, y, -z+1/2$ ; (v)  $-x, y+1, -z+1/2$ ; (vi)  $x, y-1, z$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

| $D-H\cdots A$           | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------|-------|-------------|-------------|---------------|
| O14B—H14B $\cdots$ O14A | 1.21  | 1.25        | 2.462 (10)  | 180           |
| C11B—H11B $\cdots$ O13B | 0.95  | 2.49        | 2.830 (14)  | 101           |